



# POLICY

## PREVENTING HEAT ILLNESS IN SPORT

**The object of this document is to prevent injury, and possible death, from heat illness in sport and activity by assisting officials, coaches and participants to recognise and manage potentially dangerous heat situations.**

**This is achieved by:**

1. Alerting sporting bodies and participants of the risk of heat illness from physical activity in hot weather conditions.
2. Educating sporting bodies and participants on methods of minimising the risk of heat illness and the avoidance of situations that may worsen heat illness.

# HOT WEATHER POLICY FOR THE GENERAL POPULATION

## ***DISCUSSION***

The risk of heat illness from vigorous exercise or high intensity sport is significant. It can range from cramps, through heat exhaustion to heat stroke, coma, and death (Mitchell 1994).

During competition, competitors may produce 15 - 20 times the heat they produce at rest. Dissipation of this excess heat is primarily achieved through sweating. If the body's ability to dissipate heat is compromised, core temperature in an average size individual may rise by one degree Celsius for every five minutes of exercise if no temperature-regulating mechanisms are activated (Nadel 1977). If an individual's core temperature is above 40 degrees Celsius (normal 37 degrees) the risk of heat injury is significant. Rectal temperature higher than 41 degrees Celsius is dangerous.

Factors which impair the body's ability to dissipate heat are:

1. high ambient temperature;
2. solar radiation;
3. humidity (which compromises the efficacy of sweating); and
4. dehydration.

These factors significantly increase the risk of heat illness occurring.

Sports heat illness can occur with high intensity exercise in cool conditions and with well-hydrated participants.

Because sports heat stress is complex, and because individual responses to heat stress vary, it is not possible to provide overall recommendations about limiting conditions to cover all sports. Since heat stress varies with exercise intensity, potential for heat illness may be categorised according to the exercise characteristics of the sport. The following sports involve reducing levels of exercise intensity and therefore reducing levels of risk.

1. endurance running in competition or training (higher intensity/higher risk);
2. football codes and hockey;
3. tennis; and
4. cricket (lower intensity/lower risk).

## TEMPERATURE

The tables below provide approximate guides to weather conditions and appropriate individual and organisational responses. Although temperature ranges are given, there are not clear demarcations in risk between ranges.

### **Ambient Temperature**

Easily understood, most useful on hot, dry days

<b>Ambient temperature</b>	<b>Relative humidity</b>	<b>Risk of thermal injury</b>	<b>Possible modifying action for vigorous sustained activity</b>
15 - 20		Low	Heat illness can occur in distance running. Caution over-motivation.
21 - 25	< 60%	Low - moderate	Increase vigilance. Caution over-motivation.
26 - 30	< 50%	Moderate	Moderate early pre-season training intensity. Reduce intensity and duration of play/training; take more breaks.
31 - 35	< 30%	High - very high	Limit intensity. Limit duration to less than 60 minutes per session.
36 and above	< 25%	Extreme	Consider postponement to a cooler part of the day or cancellation.

### **WBGT**

Further guidance might be gained from the Wet Bulb Globe Temperature (WBGT) index. The WBGT is particularly useful for hot, humid days.

<b>WBGT</b>	<b>Risk of thermal injury</b>	<b>Possible modifying action for vigorous sustained activity</b>
< 20	Low	Heat illness can occur in distance running. Caution over-motivation.
21 - 25	Moderate to high	Increase vigilance. Caution over-motivation. Moderate early pre-season training intensity and duration. Take more breaks.
26 - 29	High - Very high	Limit intensity. Limit duration to less than 60 minutes per session.
30 and above	Extreme	Consider postponement to a cooler part of the day or cancellation (allow swimming).

## **STRATEGIES FOR REDUCING THE RISK OF HEAT ILLNESS (General Population)**

◆PO Box 237, Dickson ACT 2602◆  
 ◆Telephone (02) 6230 4650◆Facsimile (02) 6230 5908◆  
 ◆E-mail smanat@sma.org.au◆Web www.sma.org.au

The following strategies are intended for the general population that does not fall into any of the listed 'At Risk' categories. 'At Risk' participants should consult the recommendations for their particular population sector.

### *1. Timing of games*

Games and sporting activities involving moderate to high intensity exercise should be scheduled to avoid conditions where ambient temperature exceeds or is likely to exceed 36 degrees Celsius or where WBGT exceeds 30 degrees Celsius.

In most parts of Australia players are likely to be exposed to their highest risk of heat injury in the months of December, January and February, although in some regions this level of risk extends into March and April. This is in part due to high ambient temperatures that are prevalent during this period, and lack of match fitness of players participating in traditional winter sports such as Australian Rules Football.

Where possible, especially in January and February, games should be scheduled to start before 9 am. or after 6 pm. Early morning or night games minimise the risk of encountering unacceptable conditions at these times of year. This is especially so where these games are to be played in a locations with a history of relatively high WBGT.

### *2. Acclimatisation*

If games or activities are to be conducted after long periods of cooler conditions, participants should strive to be fully acclimatised prior to participation.

Physiological adaptations to exercising in the heat are rapid and can occur after 3-5 days in a hot environment. Full acclimatisation can take 10-14 days or longer. The initial response is an expansion of the plasma volume; then, over several days, this returns to normal and the sweat rate increases with sweating starting earlier and a more dilute sweat being produced.

There is evidence that exercising in sweat clothing to the point where heat strain is induced can give some degree of acclimatisation (Dawson et al). The training must induce heat strain over several days, and care must be taken that adequate hydration occurs during these training sessions.

Doing some form of submaximal exercise in a heat chamber will also give some degree of acclimatisation; but its practicality in a team sport, except possibly in individual cases, is limited.

Some level of acclimatisation will occur in players coming out of summer. This, however, is usually countered by the lack of match fitness in athletes at this time of year.

What can be done easily is to educate athletes to train themselves to play and train with copious fluids already on-board. Further it must be emphasised to the players that they

MUST consume fluids containing 6%-8% carbohydrate - in warm/hot conditions, muscle glycogen utilisation is much higher. (Febbraio 1992). The consumption of carbohydrate containing fluids has been proven to improve performance in the heat and, more importantly, delay the onset of exercise-induced heat exhaustion (Febbraio 1992, Davies et al 1988) and, hence, probably help prevent heat stroke.

### *3. Hydration*

The more athletes sweat, the more fluid they must consume to avoid dehydration. High levels of dehydration may increase the risk of heat stress. To diminish the risk of heat stress fluid should be consumed before, during and after activity.

It is recommended participants drink at least 7-8 ml of fluid per kg of body mass (average is about 2 cups) no more than 2 hours before exercising to promote adequate hydration and allow time for excretion of excess water.

During exercise it is recommended that participants should drink fluid at regular intervals to replace water lost through sweating. Participants should aim to drink at least 3 ml per kg of body mass (about 250 ml for the average athlete of around 70 kilograms every 15 to 20 minutes or 2–3 cups every hour). However, this may vary dependent on the rate of sweating. Fluid taken should be cooler than the ambient temperature.

Water is considered an adequate fluid option for activities lasting up to one hour.. Participants in events or activities exceeding one hour are recommended to use carbohydrate-based sports drinks as a means of replacing fluids, carbohydrates and electrolytes lost during prolonged activity.

In high risk conditions players should be encouraged to drink fluids at scheduled drinks breaks and should be provided convenient access to fluids during activity without unnecessary interruption to the game or event.

Officials and event organisers should also consider including additional drinks breaks for players in conditions of high risk.

In regard to post-event rehydration, it needs to be remembered that this can take 24 hours or more.

### *4. Player rest and rotation*

In conditions of high risk participants should be provided opportunities to rest through the use of player interchange or substitution. The period of rest should be determined by the ambient temperature and WBGT at the time of the event or activity. For ambient temperatures greater than 26 and less than 30 degrees Celsius and for WBGT temperatures greater than 21 degrees Celsius and less than 25 degrees Celsius, all players should be rested for at least 10% of the period they would normally participate.

For example, if the activity normally runs for 60 minutes, the rest period for the player should comprise at least 6 minutes during the period.

For situations where the ambient temperature is greater than 31 degrees and less than 35 degrees Celsius and the WBGT is greater than 26 degrees Celsius and less than 29 degrees Celsius, all players should be rested for at least 25% of the period in which they would normally participate.

This may be achieved by rotation of players through an interchange bench or via the reduction in the regular playing time for all players.

For events played in high risk conditions that do not have a specified playing time, players should be permitted to take rest breaks from activity equivalent to 3 minutes for every 30 minutes of activity.

The positive effects of rest breaks should also be maximised by employing the following strategies:

- allowing players to rest in naturally shaded areas or providing portable structures that create shade where and when required;
- providing fans and ice packs; and
- providing additional fluids to allow participants to spray or douse themselves to assist cooling.

## *5. Pre-cooling*

Pre-cooling by cool water immersion or the wearing of ice vests has been demonstrated to increase athletic performance in endurance sports. This practice could be of benefit to many athletes. However, it must be noted that the effects of a pre-cooling manoeuvre are reduced rapidly by a warm up. Therefore, any pre-cooling strategy must be undertaken in concert with a vastly reduced warm-up if it is to be effective.

## *6. Clothing*

Light coloured, loose fitting clothes, of natural fibres or composite fabrics, with high wicking (absorption) properties that provide for adequate ventilation are recommended as the most appropriate clothing in the heat. This clothing should further complement the existing practices in Australia that protect the skin against permanent damage from the sun.

# CHILDREN AND HEAT

The physiological and structural difference between children and adults places children at a greater risk of suffering from heat illness. These differences impact on a child's ability to respond to environmental heat and acclimatise to heat. These differences include:

- a larger surface area/body mass ratio which affects their ability to dissipate heat when environmental temperature is greater than skin temperature (Falk 1998). This can be an advantage when heat loss is necessary, but is a disadvantage when radiant or convective heat gain occurs;
- immature sweating mechanisms which require a greater increase in body temperature before the onset of sweating (Araki et al 1979); and
- fewer and smaller sweat glands which limit the production of sweat (Araki et al 1979, Falk 1998, Wagner et al 1974).

## HOT WEATHER POLICY FOR CHILDREN

**At ambient temperature greater than or equal to 30 degrees Celsius, children have greater difficulty getting rid of heat than adults.**

## ***STRATEGIES FOR REDUCING THE RISK OF HEAT ILLNESS (Children)***

The following strategies should be considered for sport and physical activities involving children. The strategies should be considered in conjunction with strategies for reducing the risk of heat illness for the general population and the hot weather policy for children.

### *1. Shade and drinks*

Organisers of activities that are conducted in hot conditions must provide sufficient shade and regular drinking opportunities. This is particularly critical where the fitness and state of acclimatisation of the young participants are uncertain.

It is recommended that water or fluids be provided whenever children are being active. More fluid, however, appears to be consumed by young people when the drinks offered are perceived as palatable to them. Therefore, for children and adolescents having trouble drinking adequate tap water, flavoured drinks may need to be considered. Conversely, the high energy content of some flavoured drinks may be unnecessary during exercise in athletes who have a genuine rather than an aesthetic need to lower body fat levels.

It is recommended that young athletes begin regular drinking routines using water or fluids during training and competition. Regular and effective drinking practices should become habitual to young athletes before, during and after activity. Individuals should monitor weight changes before and after workouts and know the amount of fluid that they are likely to require.

## *2. Acclimatisation and overweight children*

In addition to the risks associated with activity in the heat for unfit and unacclimatised young people, coaches/supervisors of overweight children and adolescents should take extra precautions to lessen the potential for heat gain. It is recommended that, whenever activity in hot conditions is unavoidable with these children, coaches/supervisors decrease the volume and duration of physical activity and increase opportunities for drinking, rest and shade as a matter of priority.

At the onset of hot weather, the young athlete may take longer to acclimatise. It is therefore recommended that training volumes (duration and intensity) decrease during the first few weeks of hot weather. Increased times for rest, using access to shade more frequently and increasing the number of mandatory drinking breaks are recommended for the young athlete when the weather becomes noticeably hotter.

## *3. Clothing*

In addition to the clothing recommendations made for the general population, it is recommended that summer-based sporting organisations select uniforms that minimise heat gain and that coaches, teachers and parents encourage children and adolescents to wear appropriate clothing in layers that can be easily removed during activity.

## *4. Heat illness register*

To improve the understanding of activity in the heat by children and adolescents, it is recommended that a register of heat-related illness be established. This may comprise a system within which all aspects of heat-related illness incidents are recorded. Items of note may include the individuals afflicted and their symptoms, the time of the incident, the environmental conditions, the physical activity undertaken, the immediate treatment and subsequent action taken.

The system is recommended to help identify individuals who have previously experienced some form of heat illness and therefore may require additional attention to ensure that they adopt prevention strategies.



# RECOMMENDATIONS FOR THE DEVELOPMENT AND RECOMMENDATIONS FOR WAYS IN WHICH THE SPORTS INDUSTRY CAN REDUCE THE RISK OF HEAT ILLNESS

The following recommendations are provided to help identify long and short-term objectives for the sports industry so as to reduce significantly the risk of dangerous and sometimes catastrophic incidents occurring as a result of activity in hot weather.

## ***Adoption of policy for children***

All junior sporting clubs and associations or clubs and associations involving junior participants should immediately familiarise themselves with SMA's Hot Weather Guidelines and Beat the Heat brochure. These are available as a web download from [www.sma.org.au](http://www.sma.org.au) .

All junior sporting clubs or clubs involving junior participants are encouraged to purchase a dry bulb thermometer to measure ambient air temperature on-site to ensure local conditions are accurately measured.

## ***Adoption of strategies***

All sporting clubs and associations should develop or add to their existing policies or rules, the 'Strategies for Reducing the Risk of Heat Injury' detailed in SMA's Hot Weather Guidelines.

## ***Central measurement***

To overcome the current barriers for sporting clubs and associations measuring WBGT, it is recommended that clubs and associations avail themselves of the local weather service provided by the Bureau of Meteorology through SMA websites. .

## REFERENCES

1. Adolph E.E. Desert *In: Newburgh L.H. (Ed) Physiology of heat regulation and the science of clothing.* Philadelphia, W.B. Saunders, 1949.
2. American College of Sports Medicine, Prevention of heat injuries during distance running - Position Stand. *Med.J.Aust.* 1984 Dec.876
3. Armstrong, L.E. and Maresh, C.M. Exercise-heat tolerance of children and adolescents. *Pediatric Exercise Science* 7 239-252, 1995.
4. Araki, T., Y. Toda, K. Matsushita, and A. Tsujino. Age differences in sweating during muscular exercise. *Journal of Sports Medicine and Physical Fitness.* 4:239-248, 1979.
5. Bar-Or, O., Blimkie, C.J.R., Hay, J.A., MacDougall, J.D., Ward, D.S. and Wilson, W.M. Voluntary dehydration and heat intolerance in cystic fibrosis. *Lancet*, 33: 696-699, 1992.
6. Bar-Or, O., Dothan, R., Inbar, O., Rothstein, A. and Zonder, H. Voluntary hypohydration in 10- to 12-year old boys. *Journal of Applied Physiology*, 48: 104-108, 1980.
7. Bar-Or, O. and Wilk, B. Water and Electrolyte Replenishment in the Exercising Child. *International Journal of Sport Nutrition*, 6: 93-99, 1996.
8. Bolster D.R., Trappe SW, Short KR, Scheffield-Moore M, Parcell AC, Schulze KM, Costill DL. Effects of precooling on thermoregulation during subsequent exercise. *Med-Sci-Sports-Exerc.* Vol 31, No. 2, pp 251-257. 1999
9. Brooks, G.A., Fahey, T.D. and White, T.P. *Exercise Physiology: Human Bioenergetics and Its Applications* (2<sup>nd</sup> Edition) California, Mayfield Pub. Co. 1995 p. 433.
10. Chao T.C.,Sinniah R.,Pakiam J.E. Acute heat stroke deaths. *Pathology* 1981 13:145-156
11. Clark, R.P. and Edholm O.G. *Man and his heat environment.* London, Edward Arnold Publishers, 1985.
12. Coyle, E.F. and Montain, S.J. Carbohydrate and fluid ingestion during exercise: Are there trade-offs? *Medicine and Science in Sports and Exercise*, 24: 671-678, 1992.
13. Davies J.M.,Burgess W.A.,Slenz C.A.,Bartoli W.P.,Pate R.R. Effects of ingesting 6% and 12% glucose/electrolyte beverages during prolonged intermittent cycling in the heat. *Eur.J.Appl.Physiol.* 57:563-569,1988
14. Dawson B. Exercise Training in Sweat Clothing in Cool Conditions to Improve Heat Tolerance. *Sports- Med.*17(4):233-244,1994
15. Drinkwater, B.L., I.C. Kupprat, JE. Denton, J.L. Crist, and S.M. Horvath. Response of prepubertal girls and college women to work in the heat. *Journal of Applied Physiology* 43(6):1046-1053, 1977
16. Engeli D.B.,Maller O.,Sawka M.N.,Francesconi R.N.,Drolet L Young A.J. Thirst and fluid intake following graded hypohydration levels in humans. *Physiol-Behav.* 1987; 40(2):229-36

17. Epstein Y, Moran DS, Shapiro Y, Sohar E, Shemer J. Exertional heat stroke, a case series. *Med.Sci.Sports.Exerc.*, Vol.31, No.2, pp224-228, 1999
18. Falk, B., Physiological and health aspects of exercise in hot and cold climates. In: *The Child and Adolescent Athlete*, O. Bar-Or (Ed.). Oxford, London: Blackwell Science Ltd, 1996, pp. 326-349
19. Falk, B. Effects of heat stress during rest and exercise in the paediatric population. *Sports Medicine* 25(4): 221-240, 1998.
20. Falk, B., Bar-Or, O. and MacDougall, J.D. Thermoregulatory responses of pre- mid- and late-pubertal boys to exercise in dry heat. *Medicine and Science in Sports and Exercise*. 24(5): 688-694, 1992
21. Febbraio M.A. Heat Stress as a factor in AFL football. *AFL Grand Final symposium notes* 1992
22. Febbraio M.A., Snow R.J., Hargreaves M., Stathis C.G., Martin I.K., Carey M.F. Muscle metabolism during exercise and heat stress: effect of acclimatisation. *J.Appl.Physiol.* 76(2):589-597, 1994.
23. Gisolfi C.V., Duchman S.M. Guidelines for optimal replacement beverages for different athletic events. *Med-Sci-Sports-Exerc.* 1992 Jun; 24(6): 679-87
24. Gopinathan P.M., Pichan G., Sharma V.M. Role of dehydration in heat stress - induced variations on mental performance. *Arch-Environ-Health*. 1988 Jan-Feb; 43(i):15-17
25. Gutierrez G. Solar injury and heat illness, treatment and prevention in children. *Phys Sports Med* 1991;19(5);67-80
26. Hales J.R.S., Richards D.A.B. *Heat stress:Physical exertion and environment*. Elsevier Science Publishers, 1987
27. Kenny WL. The Older Athlete, exercise in hot environments. *Sports Science Exchange*, No.44, Vol.6,
28. Iuliano, S., Naughton, G., Collier, G. and Carlson, J.S. Examination of the self-selected fluid intake practices by junior athletes during a simulated duathlon event. *International Journal of Sports Nutrition*, 8: 10-23, 1998
29. Knochel J.P. Dog days and siriasis. How to kill a football player. *JAMA* 1975 23:513-515
30. Krivosic, Horber R., Adnet P., Reyford H. Relationship between exercise-induced myolysis and malignant hyperthermia. *Br-J-Anaesth*. 1991 Aug; 67(2): 221
31. Lyons T.P., Riedesel M.L., Meuli L.E., Chick T.W. Effects of glycerol-induced hyperhydration prior to exercise in the heat on sweating and core temperature. *Med.Sci.Sports.Exerc.*, Aug 1990, 477-483
32. Malina, R.M. and Bouchard, C. *Growth, Maturation and Physical Activity*. Champaign, Il., Human Kinetics, 1991.
33. Maxwell NS, Gardner F, Nimmo M. Intermittent running: muscle metabolism in the heat and effect of hypohydration. *Med.Sci.Sports.Exerc.*, Vol.31, No.5, pp 675-683, 1999
34. Meir R.A., Davie A.J. & Ohmsen P. Thermoregulatory responses of Rugby League Footballers playing in warm humid conditions, *Sport Health*, 1990;8(4):11-14.
35. Meyer, F., Bar-Or, O., MacDougall, J.D. and Heigenhauser, G.J.F. Drink composition and the electrolyte balance of children exercising in the heat. *Medicine and Science in Sports and Exercise*, 27: 882-887, 1995.

36. Meyer, F., Bar-Or, O., Salsberg, A. and Passe, D. Hypohydration during exercise in children: Effect on thirst, drink preferences and rehydration. *International Journal of Sports Nutrition*, 4: 22-35, 1994.
37. Mitchell B.H. (1990)The effects of dehydration and heat stress as applied to Collingwood Footballers. (*Unpublished*)
38. Mitchell BH. AFLMOA position paper on the prevention of heat injuries in football.*Sport Health*.Vol.12,No.4,pp10-18.Dec 1994
39. Mitchell BH.Prevention of dehydration and heat injuries in football.*Sports Coach*. Vol.18, No.2.pp10-13.Winter 1995
40. Pascoe, D., Bellingar, T.A., and McClusky, B.S. Clothing and Exercise: Influence of clothing during exercise / work in environmental extremes. *Sports Medicine* 18(2): 94-108, 1994.
41. Pohl A.P.,O'Halloran M.W.,Pannall P.R. Biochemical and physiological changes in football players. *Med.J.Aust.* 1: 467-470,1981
42. Pyke F.S. & Hahn A.G. Body Temperature Regulation in Summer Football,*Sports Coach*,1980;4(3):41-43.
43. Rasch W.,Samson P.,Cote J.,Cabanac M. Heat loss from the human head during exercise. *J-Appl-Physiol.* 1991 Aug; 71(2):590-5
44. Raven P.B. Is cardiac filling pressure the limiting factor in adjusting to heat stress. *Yale-J-Biol-Med.*,1986 May-June; 59(3):267-79
45. Roberts WO. Must the race go on? *Phys Sports Med* 1995;23(7):3
46. Savdie E.,Prevedoros H.,Irish A.,Vickers C.,Concannon A., Darveniza P.,Sutton J.R. Heat stroke following Rugby League football. *Med J Aust* 1991: 155;636-639
47. Sawka M.N.,Young A.J.,Francesconi R.P.,Muza S.R.,Pandolf K.B Thermoregulatory and Blood responses during exercise at graded hypohydration levels. *J.Appl.Physiol.*,1985 Nov; 59(5):1394-401
48. Sawka MN, Coyle EF. Influence of body water and blood volume on thermoregulation and exercise performance in the heat. *Exercise and sport Sciences Reviews*.Vol.27,pp167-218, 1999
49. Tek D.A.,Olshaker J.S, Hyperthermia, pulmonary oedema, and disseminated intravascular coagulation in an 18-year-old military recruit[clinical conference] *Ann-Emerg-Med.* 1990 Jun; 19(6): 715-22
50. Terrados, N. and Maughan, R.J. Exercise in the heat: Strategies to minimize the adverse effects on performance. *Journal of Sports Sciences*, 13: S55-S62, 1995.
51. Wagner, J.A., Robinson, S., Murino, R.P. Age and temperature regulation of humans in neutral and cold environments. *Journal of Applied Physiology.* 37: 562-565, 1974.
52. Wells T.D.,Jessup G.T.,Langlotz K.S. Effects of sunscreen use during exercise in the heat. *The Phys Sports Med* Vol 12,No.6,June 84
53. Young MC, Fricker PA, Thomson NJ, Lee KA. Sudden death due to ischaemic heart disease in young aboriginal sportsmen in the Northern Territory, 1982-1996. *Med J Aust* 1999 May 3;170(9):425-8